

BEVEL UP IS OPEN SOURCE. DOWNLOAD | PRINT | POST | SCROLL BVLUP.COM

GO WITH THE VEIN FLOW
AND INJECT TOWARD THE HEART
Always inject into a vein and never an artery.
Veins are close to the surface of the skin and springy to the touch.
Arteries are located deeper in the body and not as visible as veins.
If you hit an artery, it will hurt and your blood will look foamy.
If this happens, apply pressure to the wound and seek medical attention if the bleeding does not stop.

BEVEL DOWN
THE BEVEL IS THE SLANTED EDGE AT THE POINT OF THE SYRINGE
The only time the bevel should face down is when drawing up from a cooker.
You can protect the bevel by using a cotton filter.
100% cotton filters are best.
The bevel becomes dull after using it just once.

GLASS STRAIGHT STEM
CRACK, ROCK COCAINE
Brass screens are best. Ask for them at your SSP.
If you are using a scrubber for a screen, copper (Chore Boy) is better than steel wool, which can break apart and burn your mouth and throat when it gets hot.
Pre-burn your scrubber to release toxic fumes.
If sharing a pipe, use your own rubber mouthpiece (sparkplug cover or rubber bands).

CHASING WITH FOIL
ROCK/CRACK FENTANYL METH
Once the substance has liquefied, keep it moving by tilting your foil.
Use a glass stem or a straw made of foil to chase the smoke.
Take it slow, especially if it's your first time smoking or you haven't smoked in a while.
Remember that it's still possible to overdose when smoking, so give yourself a few minutes to see how you feel after that first inhale.

WATER DRIPPING
Water dripping is more effective than sniffing (liquid helps absorption).
Crush drugs as much as possible to reduce irritation.
Water from best to worst: saline, sterile, distilled, bottle or tap.
For a weaker solution, add more water.
Prep small amounts—storing mixtures can grow bacteria or mold. This is extremely dangerous.
Break the point off a syringe (dispose of it safely) or use a dropper.
P.S. This only works for drugs that dissolve in water like fentanyl, heroin, meth, and powder coke. Water dripping does not work for crack or pills.

TRY BOTH ARMS
BEFORE GOING IN HANDS, WRISTS, FEET, GROIN AND NECK
Arms usually have three accessible veins.
To find a vein, try using a soft tie or tourniquet.
Use a slip knot so that you can easily pull the tie off after injecting.
If possible, release the tie after your register (flag), but before you push in your hit.
If you are afraid you will miss your hit when you release the tie, push your hit in slow.
Ask your SSP for tourniquets and other supplies.

BEVEL UP
Position the syringe with the bevel facing up so the sharp tip can pierce the skin first, easing the way for the rest of the needle.

GLASS BUBBLE PIPE
METH, FENTANYL
Inhale slow and exhale fast.
Holding in your hit won't get you higher but it will damage your lungs.
Heat the pipe from underneath and keep the flame moving around the bowl so you melt the drugs instead of burning them.
This helps prevent the pipe from cracking and wastes less drugs.

CHASING WITH FOIL
ROCK/CRACK FENTANYL METH
Prepare a foil rectangle that is about 6" x 5" (the length and width of your hand).
If the foil has a shiny, plastic coating, pre-burn the foil by running the shiny side over a flame.
With the dull side facing down, place drugs on the foil and heat from underneath, keeping the lighter a few inches below the foil.

BOOTY BUMPING
Mix drugs with 1-2 ml of clean water.
Make sure drugs have dissolved completely—heating in a cooker can help.
Break the point off a syringe (dispose of it safely) or use a lube applicator.
Lube applicators are more effective because they are designed to deliver fluid to the rectal cavity. It also holds more liquid than a syringe.